arXiv:quant-ph/0109093v3 1 Dec 2003

MODIFIED DE BROGLIE-BOHM APPROACH TO QUANTUM MECHANICS

Moncy V. John
Department of Physics, St. Thomas College, Kozhencherri 689641, Kerala, India.

Abstract

A modified de Broglie-Bohm (dBB) approach to quantum mechanics is presented. In this new deterministic theory, which uses complex methods in an intermediate step, the problem of zero velocity for bound states encountered in the dBB formulation does not appear. Also this approach is equivalent to standard quantum mechanics when averages of observables like position, momentum and energy are taken.

Key words: quantum theory of motion, de Broglie-Bohm approach, trajectory representation, complex methods.

PACS No(s): 03.65.Bz

1 INTRODUCTION

In the de Broglie-Bohm quantum theory of motion (dBB) [1-4], which provides a deterministic theory of motion of quantum systems, the following assumptions are made [3]:

(A1) An individual physical system comprises a wave propagating in space and time, together with a point particle which moves continuously under the guidance of the wave.

(A2) The wave is mathematically described by $\Psi(x,t)$, a solution to the Schrodinger's wave equation.

(A3) The particle motion is obtained as the solution $x(t)$ to the equation

$$\dot{x} = \frac{\hbar}{2im} \frac{\left(\Psi^\star \frac{\partial \Psi}{\partial x} - \frac{\partial \Psi^\star}{\partial x} \Psi\right)}{\Psi^\star \Psi}, \tag{1}$$

with the initial condition $x = x(0)$. An ensemble of possible motions associated with the same wave is generated by varying $x(0)$.

(A4) The probability that a particle in the ensemble lies between the points x and $x + dx$ at time t is given by $\Psi^\star \Psi \, dx$.

In spite of its success as a deterministic theory, this scheme has the difficulty that, for bound state problems, the time-independent part ψ of the wave function is real and hence the velocity of the particle turns out to be zero everywhere [3]. This feature, that the particle is always at rest, is not a satisfactory one in a quantum theory of motion.

To have a better understanding of this problem, let us recall that in the the classical Hamilton-Jacobi theory, the trajectory for a particle may be obtained from the equation

$$\frac{\partial S}{\partial t} + H\left(x, \frac{\partial S}{\partial x}\right) = 0 \tag{2}$$

by first attempting to solve for the Hamilton-Jacobi function $S = W - Et$ and then integrating the equation of motion

$$p = m\dot{x} = \frac{\partial S}{\partial x} \tag{3}$$

with respect to time. However, since the classical Hamilton-Jacobi equation is not adequate to describe the micro world, one resorts to the Schrodinger equation $i\hbar \partial\Psi/\partial t = -(\hbar^2/2m)\partial^2\Psi/\partial x^2 + V\Psi$ and solves for the wave function Ψ. The basic idea of dBB is to put Ψ in the polar form $Re^{iS/\hbar}$ and then to identify S as the Hamilton-Jacobi function. This gives us Eq. (1), when the particle velocity is given by Eq. (3). For wave functions whose space part is real, the Hamilton's characteristic function $W = constant$ and hence the velocity field is zero everywhere, which is the difficulty mentioned above. This is certainly an undesirable feature in an ontological interpretation of quantum mechanics, which claims to make the theory intelligible and causal [2]. It is true that dBB is altogether a non-Copenhagen physical theory which requires a 'quantum intuition' and it is not justifiable to criticize dBB for its failure to return to classical mechanics. However, the fact that the theory dubs the material system in the most important and wide class of bound state problems as stationary, irrespective of its position and energy, is counter-intuitive and hence the theory cannot be termed as quantum theory of motion, at least in these cases.

Another deterministic approach to quantum mechanics, which also claims the absence of this problem, is the Floyd-Faraggi-Matone (FFM) trajectory representation developed by Floyd [5]. In one dimension, Floyd uses the generalized Hamilton-Jacobi equation

$$\frac{(W')^2}{2m} + V - E = -\frac{\hbar^2}{4m}\left[\frac{W'''}{W'} - \frac{3}{2}\left(\frac{W''}{W'}\right)^2\right], \tag{4}$$

where W' ($'$ denotes $\frac{\partial}{\partial x}$) is the momentum conjugate to x. Recently Faraggi and Matone [6] have independently generated the same equation (referred to as the quantum stationary Hamilton-Jacobi equation) from an equivalence principle free from any axioms. Floyd argues that the conjugate momentum W' is not the mechanical momentum; instead, $m\dot{x} = m\partial E/\partial W'$. The equation of motion in the domain $[x, t]$ is rendered by the Jacobi's theorem. For stationarity, the equation of motion for the trajectory time t, relative to its constant coordinate τ, is given as a function of x by [5, 6, 7]

$$t - \tau = \frac{\partial W}{\partial E}, \tag{5}$$

where τ specifies the epoch. Thus the dBB and FFM trajectory representations differ significantly in the use of the equation of motion, though they are based on equivalent generalized Hamilton-Jacobi equations. The FFM trajectory representation does not claim equivalence with standard quantum mechanics in the predictions of all observed phenomena.

In this letter, we present a different and modified version of the dBB that surmounts the problem mentioned earlier. In addition, its formulation is closer to the classical Hamilton-Jacobi theory than even the conventional dBB. We apply the scheme to a few simple problems and find that it is capable of providing a deeper insight into the quantum phenomena. Also the new scheme is constructed in such a way that it is equivalent to standard quantum mechanics when averages of observables like position, momentum and energy are taken.

In an intermediate step in the new formulation, we consider the analytic continuation of the spacetime into complex, producing a complex spacetime. It is well known that such complex spacetimes are put to remarkable use in the theory of relativity. For example, the transformation of one solution of Einstein equation into another can on many occasions be accomplished by means of a simple complex switch of coordinates. Open and closed Friedmann models, de Sitter and ant-de Sitter spacetimes, Kerr and Schwarschild metrics etc. are related by complex substitutions. The use of complex variables in relativity extends to more sophisticated ones like spin coefficient formalism, Ashtekar formalism, Twistor theory etc. Recently, it was demonstrated that a Friedmann model with complex scale factor leads to a remarkably good nonsingular cosmological model, with none of the main cosmological problems in standard cosmology appearing in it [8]. The Wick rotation of the time coordinate t in Minkowski space to $i\,t$, which changes its metric into Euclidean or reversing, going from Euclidean to Minkowski metric is already familiar in quantum field theory and quantum gravity as Euclidean Feynman path integration. Even in quantum mechanics, the technique of considering the space variable as complex is familiar [9]. Some more specific usage of complex methods include the interpretation of spin angular momentum as orbital momentum but taken about a complex center of mass and the interpretation of magnetic moment as the electric dipole moment taken about a complex center of charge [10]. The present modified dBB scheme may be considered as yet another application of complex methods in quantum theory.

The paper is organized as follows. In Sec. 2, we present the basic formalism and in Sec. 3, we show that the new scheme is constructed to give the same statistical predictions as that in standard quantum mechanics. In section 4, we apply the scheme to a few simple problems while section 5 deals with the problem of action variable in the new scheme. The last section gives our conclusions and discussion.

2 THE GENERAL FORMALISM

We first note that in the dBB, the substitution $\Psi = Re^{iS/\hbar}$ in the Schrodinger equation gives rise to two coupled partial differential equations, one of which is a conservation equation for probability density, and this leads to a situation apparently different from the Hamilton-Jacobi theory for individual particles. On the other hand, we note that a change of variable $\Psi \equiv \mathcal{N} e^{i\hat{S}/\hbar}$ in the Schrodinger equation (where $\mathcal{N}$ is a normalisation constant) gives a quantum-mechanical Hamilton-Jacobi equation [11]

$$\frac{\partial \hat{S}}{\partial t} + \left[\frac{1}{2m} \left(\frac{\partial \hat{S}}{\partial x} \right)^2 + V \right] = \frac{i\hbar}{2m} \frac{\partial^2 \hat{S}}{\partial x^2}, \tag{6}$$

which is closer to its classical counterpart and which also has the correct classical limit. Considering $\hat{S}$ as a modified Hamilton-Jacobi function brings the expression (3) for the conjugate momentum to the form

$$m\dot{x} \equiv \frac{\partial \hat{S}}{\partial x} = \frac{\hbar}{i} \frac{1}{\Psi} \frac{\partial \Psi}{\partial x}. \tag{7}$$

In this paper, we attempt to modify the dBB scheme in a manner similar to the latter approach; i.e., we retain assumptions A1 and A2 whereas in A3, we identify $\Psi \equiv \mathcal{N} e^{i\hat{S}/\hbar}$ and use the expression (7) as the equation of motion.

Note that in this case, with Ψ as the standard solution to the Schrodinger equation in any given problem, $\dot{x}$ turns out to be a complex variable. Consequently, also x should be complex. As mentioned in the introduction, considering x as a complex variable is not entirely new, even in quantum mechanics. While discussing the quasi-classical problems, Landau and Lifshitz [9] use the technique of 'complex classical paths' extensively, as an intermediate step. We propose to consider x as complex in the general formalism of quantum mechanics, in the context of dBB theory. It is clear that when x is a complex variable, the Schrodinger equation itself becomes a complex differential equation. In the following, we consider only those solutions to such equation, obtained by replacing the real x with $x \equiv x_r + ix_i$ in the standard solution. From here onwards, we consider x as a complex variable.

We restrict ourselves to single particles in one dimension for simplicity. The formal procedure adopted is as follows: Identifying the standard solution Ψ as equivalent to $\mathcal{N} e^{i\hat{S}/\hbar}$, Eq.(7) helps us to obtain the expression for the velocity field $\dot{x}$ in the new scheme. This, in turn, is integrated with respect to time to obtain the complex path $x(t)$, with an initial value $x(0)$. This complex solution is obviously quite different from the corresponding dBB solution. The connection with the real physical world is established by postulating that the physical coordinate or trajectory of the particle is the real part $x_r(t)$ of its complex path $x(t)$. (It is worth recalling that this procedure is commonly adopted in physical problems which involve differential equations with complex solutions, as in the case of a classical harmonic oscillator.) It can be seen that in the general case, also this modified dBB trajectory $x_r(t)$ is different from the dBB trajectory.

The real part of the velocity field $\dot{x}_r$, at any point x in the complex plane may be found from Eq. (7) as

$$\dot{x}_r = \frac{\hbar}{2im} \frac{\left[\Psi^\star \frac{\partial \Psi}{\partial x} - (\frac{\partial \Psi}{\partial x})^\star \Psi\right]}{\Psi^\star \Psi}, \tag{8}$$

an expression quite analogous to the expression for the velocity field in the dBB scheme (but not identical with it, since $\dot{x}_r$ is now defined over the entire x-plane). One can also write $\dot{x}_r$ as

$$\dot{x}_r = \frac{\hbar}{2im} \frac{\left[\Psi^\star \frac{\partial \Psi}{\partial x_r} - (\frac{\partial \Psi}{\partial x_r})^\star \Psi\right]}{\Psi^\star \Psi}, \tag{9}$$

where use is made of the fact that for complex derivatives, $\frac{\partial \Psi}{\partial x} = \frac{\partial \Psi}{\partial x_r}$. It is easy to see that this is identical to the dBB velocity field at all points along the real axis $x = x_r$.

We shall note that the new scheme differs from the dBB also in one important aspect. Here, when the particle corresponds to some point x in the complex plane, its physical coordinate is x_r and the physical velocity is $\dot{x}_r$, evaluated at x using (8). Thus for the same physical coordinate x_r, the particle can have different physical velocities, depending on the point x through which its complex path passes. This in principle rules out the possibility of ascribing simultaneously well-defined physical position and velocity variables for the particle. The situation here is very different from the dBB in which for given physical position of the particle, also the velocity is known, as obtainable from Eq. (1).

Let us consider the case in which $\hat{S} = \hat{W} - Et$, where E and t are assumed to be real. Then the Schrodinger equation gives us an expression for the energy of the particle

$$E = \frac{1}{2}m\dot{x}^2 + V(x) + \frac{\hbar}{2i}\frac{\partial \dot{x}}{\partial x}. \tag{10}$$

The last term resembles the quantum potential in the dBB theory.

3 EXPECTATION VALUES

We recall that in this scheme, for the same quantum state the particle can have any number of complex paths, depending on the initial value $x(0)$. When we calculate the expectation values of dynamical variables, it should be such that this is not done for a single particle which belongs to a particular complex path, but over an ensemble of particles which corresponds to all possible complex paths. We modify assumption (A4) by postulating that the average of an observable O is obtained using the measure $\Psi^\star\Psi$ as

$$< O >= \int_{-\infty}^{\infty} O\Psi^\star\Psi \; dx, \tag{11}$$

where the integral is taken along the real axis. This can be explicitly written for x, p and E as

$$< x >= \int x\Psi^\star\Psi \; dx, \tag{12}$$

$$< p >= \int p\Psi^\star\Psi \; dx = \int \Psi^\star \frac{\hbar}{i}\frac{\partial \Psi}{\partial x} \; dx \tag{13}$$

and

$$< E >= \int E\Psi^\star\Psi \; dx = \int \Psi^\star \left(\frac{-\hbar^2}{2m}\frac{\partial^2}{\partial x^2} + V(x) \right) \Psi \; dx, \tag{14}$$

where use is made of equations (7) and (10) in equations (13) and (14), respectively. Note that we did not need to make the conventional operator replacements in the above expressions. Since the integral is taken along the real axis, these coincide with the corresponding quantum mechanical expectation values. Thus the new scheme is equivalent to standard quantum mechanics when averages of observables like position, momentum and energy are computed using the above prescription.

4 SIMPLE APPLICATIONS

To see how the scheme works, let us consider the example of the ground state solution Ψ_0 of the Schrodinger equation for the harmonic oscillator in one dimension, the space part of which is real. We have

$$\Psi_0 = \mathcal{N}_0 e^{-\alpha^2 x^2/2} e^{-iE_0 t/\hbar}. \tag{15}$$

The velocity field in the new scheme is given by Eq. (7) as

$$\dot{x} = -\frac{\hbar}{im}\alpha^2 x, \tag{16}$$

whose solution is

$$x = A e^{i\hbar\alpha^2 t/m}. \tag{17}$$

This is an equation for a circle of radius $|A| = |x(0)|$ in the complex plane (Fig. 1). As proposed earlier, we take the real part of this expression,

$$x_r = A\cos(\hbar\alpha^2 t/m), \tag{18}$$

[where it is chosen that at $t = 0$, $x(0) \equiv A$ is real] as the physical coordinate of the particle. It shall be noted that this is the same classical solution for a harmonic oscillator of frequency $\omega_0 = \hbar\alpha^2/m$. However, the nonclassical feature here is that we have a trajectory for every initial value $x(0)$, though the energy E has the value $\frac{1}{2}\hbar\omega_0$ for all such trajectories.

We can adopt this procedure to obtain the velocity field, also for higher values of the quantum number n. For $n = 1$, we have

$$\Psi_1 = \mathcal{N}_1 e^{-\alpha^2 x^2/2}\ 2\alpha x\ e^{-iE_1 t/\hbar}, \tag{19}$$

from which

$$\dot{x} = \frac{\hbar}{im}\left(-\alpha^2 x + \frac{1}{x}\right). \tag{20}$$

The solution to this equation can be written as

$$(\alpha x - 1)(\alpha x + 1) = A e^{2i\hbar\alpha^2 t/m} \tag{21}$$

or

$$x = \frac{1}{\alpha}\sqrt{1 + Ae^{2i\hbar\alpha^2 t/m}}. \tag{22}$$

Here, the solution is a product of two circles centered about $\alpha x = \pm 1$, which is plotted in Fig. 2. The physical coordinate of the particle is again given by the real part of this expression. For $n = 2$, the solution can similarly be constructed as

$$4\alpha x\left(\alpha x - \sqrt{5/2}\right)^2\left(\alpha x + \sqrt{5/2}\right)^2 = Ae^{5i\hbar\alpha^2 t/m}. \tag{23}$$

The complex path in the x-plane is plotted in Fig. 3. For cases with $n \geq 1$, these paths deviate from the classically expected circular ones. Note that in all these cases, the velocity fields are not zero everywhere, contrary to what happens in the dBB.

Now let us apply the procedure to some other stationary states which have complex wave functions. For a plane wave, we have

$$\Psi = Ae^{ikx}e^{-iEt/\hbar}, \tag{24}$$

so that the equation of motion is $\dot{x} = \hbar k/m$ and this gives

$$x = x(0) + \frac{\hbar k}{m}t, \tag{25}$$

the classical solution for a free particle.

As another example, consider a particle with energy E approaching a potential step with height V_0, shown in Fig. 4. In region I, we have

$$\psi_I = e^{ikx} + Re^{-ikx}, \qquad k = \sqrt{2mE/\hbar^2} \tag{26}$$

and, in region II,

$$\psi_{II} = Te^{iqx}, \qquad q = \sqrt{2m(E - V_0)/\hbar^2}. \tag{27}$$

The velocity fields in the two regions are given by

$$\dot{x}_I = \frac{\hbar k}{m}\left(\frac{e^{ikx} - Re^{-ikx}}{e^{ikx} + Re^{-ikx}}\right) \tag{28}$$

and

$$\dot{x}_{II} = \frac{\hbar q}{m}, \tag{29}$$

respectively. The contours in the complex x-plane in region I, for a typical value of the reflection coefficient $r \equiv R^2 = 1/2$, are given by

$$\sqrt{2}\cos 2kx_{Ir} - e^{-2kx_{Ii}} - \frac{1}{2}e^{2kx_{Ii}} = c \tag{30}$$

(where x_{Ir} and x_{Ii} are, respectively, the real and imaginary parts of x_I), and are plotted in Fig. 5. Note that also this case is significantly different from the corresponding dBB solution.

Lastly, we consider a nonstationary wave function, which is a spreading wave packet. Here, let the propagation constant k has a Gaussian spectrum with a width $\Delta k \sim 1/\sigma$ about a mean value $\bar{k}$. The wave function is given by

$$\Psi(x,t) = \mathcal{N}\left(\frac{2\pi\sigma}{\sigma^2 + i\hbar t/m}\right)^{1/2} \exp\left[-\frac{(x - i\sigma^2\bar{k})^2}{2(\sigma^2 + i\hbar t/m)} - \frac{\sigma^2 + \bar{k}^2}{2}\right]. \tag{31}$$

The velocity field is obtained from (7) as

$$\dot{x} = -\frac{\hbar}{im}\left(\frac{x - i\sigma^2\bar{k}}{\sigma^2 + i\hbar t/m}\right) \tag{32}$$

Integrating this expression, we get

$$x(t) = x(0) + \frac{\hbar\bar{k}}{m}t + i\frac{\hbar}{m}\frac{x(0)}{\sigma^2}t. \tag{33}$$

Separating the real and imaginary parts of this equation (and assuming $\bar{k}$ is real), we get,

$$x_r(t) = x_r(0) + \frac{\hbar\bar{k}}{m}t + \frac{\hbar}{m}\frac{x_i(0)}{\sigma^2}t, \tag{34}$$

$$x_i(t) = x_i(0) + \frac{\hbar}{m}\frac{x_r(0)}{\sigma^2}t. \tag{35}$$

For the particle with $x(0) = 0$, we obtain $x_r(t) = (\hbar\bar{k}/m)t$ and $x_i(t) = 0$; i.e., this particle remains at the center of the wave packet. Other particles assume different values for $x(t)$ at time t as given by the above expression, which indicates the spread of the wave packet.

5 ACTION VARIABLE

It is now of interest to calculate the action variable J for the bound state orbits in the harmonic oscillator problem considered above. If we define $J \equiv \oint m\dot{x}\ dx$, where $m\dot{x}$ is given by Eq. (7), it can be verified that $J = nh$ $(n = 0, 1, 2, 3..)$ for those particles whose complex paths enclose all the singular points of the integrand. In the general case, we shall write

$$J = \frac{\hbar}{i} \oint \frac{\partial\Psi/\partial x}{\Psi}\, dx$$

We may note that $\int \frac{\partial\Psi/\partial x}{\Psi}\, dx = \ln\Psi + \text{constant}$ has a branch cut whenever Ψ goes from positive to zero through negative. In the harmonic oscillator problem, the integral reduces to

$$J = \frac{\hbar}{i} \oint \left[-\alpha^2 x + \frac{H_n'(\alpha x)}{H_n(\alpha x)} \right] dx = \frac{\hbar}{i} \oint \frac{H_n'(\alpha x)}{H_n(\alpha x)} dx$$

where the prime denotes differentiation with respect to x. It is easily verified that for $n = 0$, J is always zero. For $n = 1$, $J = \frac{\hbar}{i} \oint \frac{1}{x}\, dx = h$ when the integration is performed over complex paths enclosing the pole $x = 0$. From Fig. 2, it is seen that this happens for the larger nests with $A > 1$ whereas for the subnests with $A < 1$, $J = 0$. (In fact, the subnests appear to be physically nonviable single particle solutions.) Similarly for $n = 2$, $J = 2h$ when the complex path encloses the poles at $x = \pm\frac{1}{\sqrt{2}\alpha}$ and so on. In all cases, J vanishes for the subnests.

Comparing the value of the action variable for the present case with that of dBB and FFM reveals the fact that the new scheme is quite different from the other representations. In the dBB, $J = 0$ by precept. FFM has shown that the J-quantization by Milne [12] can be generalized to allow microstates. The integrand, the location of poles and the branch cuts in the present case differ with that of Milne. Also note that in Milne and FFM, $J = nh$ for $n = 1, 2, 3, ..$ whereas in the new scheme $J = nh$, $n = 0, 1, 2, 3, ..$ only for the larger nests and $J = 0$ otherwise.

Having obtained the value of the action variable, one may check whether

$\partial J/\partial E = \tau$ is the orbital period in this scheme. For the harmonic oscillator, this is true since E is linear in n. Whether τ is the orbital period for the general potential is an interesting open question. Following the arguments in [11], one can see that in spite of the complexification involved, $\tau \equiv 1/\nu$ is the orbital period if the equation of motion for $w \equiv \partial \hat{W}/\partial E$ is given by $\dot{w} = \partial H(J)/\partial J$. We need to solve more examples to verify this.

Under these circumstances, one may conclude that the issue of action variable is not satisfactorily resolved for the new scheme.

6 DISCUSSION

In summary, we find that the dBB identification $\Psi = Re^{iS/\hbar}$, which itself is quite arbitrary, does not help to utilize all the information contained in the wave function while solving the equation of motion $m\dot{x} = \partial S/\partial x$. In the present formulation we use a quantum Hamilton-Jacobi equation (obtained by substituting $\Psi = \mathcal{N} e^{i\hat{S}/\hbar}$ in the Schrodinger equation), which is closer to the classical one. This formalism is already used by some authors (for example, see [13]) for other purposes. Employing the equation of state $m\dot{x} = \partial \hat{S}/\partial x$ brings the motion to a complex x-plane. The physical coordinate or trajectory of the particle is postulated to be the real part of $x(t)$. Solving the equation of motion in some bound state problems shows why the dBB scheme gives zero velocity for particles in these cases; the complex paths cross the real axis parallel to the imaginary one, so that $\dot{x}_r = 0$ at these points. The positive results we obtained for the harmonic oscillator and potential step problems themselves are indicative of the deep insight obtainable in such problems by the use of the modified dBB scheme.

It shall be noted that some of the differences between FFM and dBB trajectories are still there between FFM and modified dBB trajectories. But as claimed in the paper, modified dBB has many advantages over dBB trajectories. An outline of such differences between FFM, dBB and modified dBB trajectories can be made as follows:

(i) dBB and modified dBB need probability to obtain expectation values of observable physical quantities. But FFM trajectories do not involve probability.

(ii) dBB and modified dBB have similar equations of motion. FFM uses a different equation of motion.

(iii) dBB asserts that the possible trajectories for a particular particle should not cross. But FFM and modified dBB trajectories can cross.

(iv) For bound states, the particle in the dBB scheme does not move. But in FFM and modified dBB approaches, the particles in bound states have well-defined trajectories.

(v) The ultimate test of these trajectories lies in experiments involving time sequence of single systems (which can be carried out nowadays), rather than those involving large number of particles. FFM has already proposed some experiments which involve predictions different from standard quantum mechanics. In that sense, FFM is a testable theory. The present modified dBB formulation is at a disadvantage that as yet it does not provide a physical example where such predictions arise. An important avenue to search for in this alternative approach is a solution to this shortcoming.

A theoretical understanding of the correctness of the equation of motion used here, in comparison with the equation of motion used in FFM is highly desirable. This is particularly so since we cannot expect both theories to be true while they differ. This comparison can perhaps be made by analysing the time concept (Floydian time or some standard time) and could be clarified therefrom. Also this important question is not addressed in this paper.

The prescription of $\dot{x}$ by Eq. (7) is reminiscent of Schrodinger's seminal work [14] on the wave function. Schrodinger attempted to derive a wave equation by accepting the classical Hamilton-Jacobi equation as a phenomenological equation and by replacing $\partial W/\partial x$ in this equation with $(K/\Psi)(\partial\Psi/\partial x)$ where K is a constant. The present work can be considered to be a reversal of Schrodinger's process, in which the attempt is to obtain an equation of motion for the particle by accepting the Schrodinger wave equation as a phenomenological equation and by equating $(\hbar/i\Psi)(\partial\Psi/\partial x)$ with $m\dot{x}$. In this sense, it is closer to Schrodinger's than to dBB.

Generalization of the formalism to more than one dimension is not attempted in this letter. Its application to those other physical problems of interest shall be addressed in future publications.

Acknowledgements

Valuable comments made by Drs. Edward R. Floyd, A. Bouda, M. Matone, R. Brown and R. Carroll are acknowledged with thanks. It is a pleasure to thank Professor K. Babu Joseph, Sajith and Satheesh for encouragement and discussion.

References

[1] L. de Broglie, *J. Phys. Rad., 6^{e} serie, t.* **8**, 225 (1927); D. Bohm, *Phys. Rev.* **85**, 166, (1952).

[2] D. Bohm and B.J. Hiley, *The Undivided Universe* (Routledge, London and New York, 1993).

[3] P. Holland, *The Quantum Theory of Motion* (Cambridge University Press, Cambridge, England, 1993).

[4] R. Carroll, *Quantum Theory, Deformation, and Integrability* (North Holland, 2000).

[5] E. R. Floyd, *Phys. Rev.* **D 26**, 1339 (1982); **25**, 1547 (1982); **29**,1842 (1984); **34**, 3246 (1986); *Found. Phys. Lett.* **9**, 489 (1996); **13**, 235 (2000); *Int. J. Mod. Phys.* **A 14**, 1111 (1999).

[6] A. Faraggi and M. Matone, *Phys. Lett.* **B 450**, 34 (1999); **437**, 369 (1998); **445**, 77 (1999); *Int. J. Mod. Phys.* **A 15**, 1869 (2000).

[7] R. Carroll, *J. Can. Phys.* **77**, 319 (1999); A. Bouda, *Found. Phys. Lett.* **14**,17 (2001), quant-ph/0004044.

[8] M. V. John and K. Babu Joseph, *Phys. Lett.* **B 387**, 466 (1996); *Class. Quantum Grav.* **14**, 1115 (1997); *Phys. Rev.* **D 61**, 87304 (2000); M. V. John, Ph.D. Thesis, Cochin University of Science and Technology, Kochi, India (2000), gr-qc/0007053; M. V. John and J. V. Narlikar, *Phys. Rev.* **D 65**, 043506 (2002).

[9] L. D. Landau and E. M. Lifshitz, *Quantum Mechanics* (Pergamon Press, Oxford, 1977).

[10] E. T. Newmann and J. Winicour, *J. Math. Phys.* **15**, 1113 (1974); E. T. Newmann, *J. Math. Phys.* **14**, 102 (1973).

[11] H. Goldstein, *Classical Mechanics* (Addison-Wesley, Reading, Massachusetts, 1980).

[12] W. E. Milne, *Phys. Rev.* **35**, 863 (1930).

[13] M. R. Brown and B. J. Hiley, quant-ph/0005026.

[14] M. Jammer, *Conceptual Development of Quantum Mechanics*, (McGraw Hill, New York, 1966); E. Schrodinger, Ann. Phys. **79**, 361 (1926)



Figure 1: Complex paths in the X-plane ($X \equiv \alpha x$) for the $n = 0$ harmonic oscillator case, where contours are plotted for $X(0) =$1, 2, 3 and 4.



Figure 2: Complex paths in the X-plane ($X \equiv \alpha x$) for the $n = 1$ harmonic oscillator case, where contours are plotted for $X(0)$ =1.2, 1.35, 1.45 and 1.55.



Figure 3: Complex paths in the X-plane ($X \equiv \alpha x$) for the $n = 2$ harmonic oscillator case, where contours are plotted for $X(0)$=1.8, 1.9, 2.0 and 2.1.



Figure 4: Potential step, with $V = 0$ for $x_r < 0$, $V = V_0$ for $x_r > 0$ and energy $E > V_0$.



Figure 5: Complex paths for particles approaching the potential step. Contours for c = -4, -3, -2, -1 and 0 are plotted.